



Figure 3. Demonstration of a functional mutation with bacterial expression assay. Mutated FECH protein expressed in *E. coli* BL21 has no residual activity. Data are shown as the percentage of the FECH activity with normal control, based on the mean \pm SD of triplicate experiments. The average activity of cell extracts from pAR-HF transformed bacteria was 10.2 nmol zinc-mesoporphyrin formed per min at 37°C.

transformed bacteria was similar to that of a mock cDNA-transformed one, indicating that the mutated FECH did not exhibit any activity (**Fig 3**).

The location of mutations in the FECH gene is highly heterogeneous and there are no mutation hot spots. The predominant type of FECH gene mutations are single-nucleotide substitutions in the coding region (42%), followed by exon skipping (34%) and small deletions and insertions (21%) (Rufenacht *et al*, 1998). This is the first instance of an amino acid insertion in the FECH protein. The transition of A to G at 4 bases from the 3' terminus of intron 4 (IVS4⁻⁴, a \rightarrow g) leads to in-frame insertion of three bases in ferrochelatase mRNA (**Fig 2B**). It is suggested that recognition of the 3' splice site a(-2)g(-1) by the splicing factor U2AF³⁵ (Wu *et al*, 1999) was disturbed due to the mutation at a(-5)g(-4). As a result of misreading the 3' acceptor splice site, aberrant splicing would occur. Aberrant splicing has been classified into four types: exon skipping, cryptic site activation, intron retention, and new site creation (Nakai and Sakamoto, 1994). Our case corresponds to the last type. The observed frequency of point mutations at position -4 within 3' splice sites was 0 among 101 different point mutations in the vicinity of mRNA splice junctions (Krawczak *et al*, 1992). To our knowledge, a mutation at position -4 that generated a novel acceptor splice site in FECH gene is rare and critical for this patient.

It is not too surprising that his mother was asymptomatic, though she was found to be heterozygous for the mutation. The reason is that EPP is thought to be inherited with low clinical penetrance and may be considered as an inherited disorder that does not strictly follow recessive or dominant rules (Gouya *et al*, 1996). Other factors such as the expression level of the wild-type allele and environmental effects may contribute to modulation of the phenotype.

This work was supported in part by a Grant-in-Aid for Scientific Research (C) to 70244241 (SY) from The Ministry of Education, Science, Sports and Culture of Japan.

Shinichi Yotsumoto, Shoko Shimada, Kenjiro Terasaki, Shigeru Taketani,[†] Keiko Kobayashi,* Tekeyori Saheki,* Tamotsu Kanzaki

Departments of Dermatology and *Biochemistry, Faculty of Medicine, Kagoshima University, Kagoshima, Japan

[†]Department of Biotechnology, Kyoto Institute of Technology, Kyoto, Japan

REFERENCES

- Gouya L, Deybach JC, Lamoril J, Da Silva V, Beaumont C, Grandchamp B, Nordmann Y: Modulation of the phenotype in dominant erythropoietic protoporphyria by a low expression of the normal ferrochelatase allele. *Am J Hum Genet* 58:292-299, 1996
- Krawczak M, Reiss J, Cooper DN: The mutational spectrum of single base-pair substitutions in mRNA splice junctions of human genes: causes and consequences. *Hum Genet* 90:41-54, 1992
- Li FM, Lim CK, Peters TJ: An HPLC assay for rat liver ferrochelatase activity. *Biomed Chromatogr* 2:164-168, 1987
- McKusick VA: *Mendelian Inheritance in Man. A Catalog of Human Genes and Genetic Disorders*. Baltimore: Johns Hopkins University Press, 1998, pp. 1566-1568
- Nakahashi Y, Taketani S, Okuda M, Inoue K, Tokunaga R: Molecular cloning and sequence analysis of cDNA encoding human ferrochelatase. *Biochem Biophys Res Comm* 173:748-755, 1990
- Nakai K, Sakamoto H: Construction of a novel database containing aberrant splicing mutations of mammalian genes. *Gene* 141:171-177, 1994
- Okuda M, Kohno H, Furukawa T, Tokunaga R, Taketani S: Overexpression in *Escherichia coli*, and one-step purification of the human recombinant ferrochelatase. *Biochim Biophys Acta* 1200:123-128, 1994
- Rufenacht UB, Gouya L, Schneider-Yin X, *et al*: Systematic analysis of molecular defects in the ferrochelatase gene from patients with erythropoietic protoporphyria. *Am J Hum Genet* 62:1341-1352, 1998
- Taketani S, Inazawa J, Nakahashi Y, Abe T, Tokunaga R: Structure of the human ferrochelatase gene. Exon/intron gene organization and location of the gene to chromosome 18. *Eur J Biochem* 205:217-222, 1992
- Todd DJ: Erythropoietic protoporphyria. *Br J Dermatol* 131:751-766, 1994
- Wu S, Romfo CM, Nilsen TW, Green MR: Functional recognition of the 3' splice site AG by the splicing factor U2AF³⁵. *Nature* 402:832-835, 1999

Mechanical Stress Induced by Hyperosmolarity

To the Editor:

We have recently read the article by Dascalu *et al* (2000) in which our work was cited and hereby wish to comment on what we believe to be inaccuracies in the interpretation of our cited article.

Manuscript received February 19, 2001; accepted for publication March 12, 2001.

Reprint requests to: Dr. Stefan Kippenberger, Department of Dermatology and Venerology, Klinikum der Johann Wolfgang Goethe Universität, Theodor-Stern-Kai 7, D-60590 Frankfurt/Main, Germany.

Several lines of evidence have supported the assumption that mechanical forces applied to human cells are able to modulate fundamental events like proliferation and differentiation (Chen *et al*, 1997). Our group has already shown that different qualities of mechanical stimuli provoke different cell responses (Görmar *et al*, 1990; Kippenberger *et al*, 1999, 2000a, b). The application of cell stretch to human keratinocytes *in vitro* amplifies the proliferation cascade, whereas mechanical pressure was shown to increase differentiation signals. In the October 2000 issue of the *Journal of Investigative Dermatology*, Dascalu *et al* subjected human keratinocytes to hyperosmotic stress and discussed the cellular responses as a

quality of mechanical stress. In particular, they observed a growth inhibition due to hyperosmotic stress and referred to data from our group (Görmär *et al*, 1990) and that of Takei *et al* (1997), which in their opinion, were contradictory. This is not the case, as both cited publications made use of different qualities of mechanical stress. In the work of Takei *et al* mechanical stretch lead to enhanced proliferation of human keratinocytes, whereas our work made use of mechanical pressure giving an increase in differentiation parameters. Both these findings show no conflict and are well in harmony with the *in vivo* situation where skin stretch leads to skin enlargement without changes in skin thickness. Examples for this functional connection are given by abdominal skin growth during pregnancy or the use of skin expanders for cosmetic surgery. On the other hand, the application of pressure to the skin organ repels proliferation and instead triggers differentiation processes (hyperkeratosis, acanthosis).

Dascalu *et al* applied hyperosmotic stress to human keratinocytes and interpreted this stimulus as being mechanically relevant without showing it. They claim that hyperosmotic conditions represent a form of mechanical pressure. This is an error in logic as hyperosmotic conditions initially lead to cell shrinkage and therefore to a decrease in intracellular pressure accompanied by reduced turgidity. In contrast, the application of mechanical pressure leads to cell compression that might also increase intracellular pressure. In conclusion, we strongly suggest investigating how hyperosmotic stress alters the cell volume in the system presented and how compensating mechanisms like regulatory volume increase (RVI) counteract osmotic shrinkage. At present, the reported data give no evidence that hyperosmotic stress represents a form of mechano-stress. In this context it is essential to

use more than just one osmotically active substance in order to delineate measured cell responses to the presence of hyperosmolarity. Therefore, it maybe worthwhile to also make use of membrane-permeable osmotic active substances like urea that would be able to substantiate the relevance of cell volume in this context.

Stefan Kippenberger, August Bernd, Roland Kaufmann
Klinikum der Johann Wolfgang Goethe Universität,
Frankfurt/Main, Germany

REFERENCES

- Chen CS, Mrksich M, Huang S, Whitesides GM, Ingber DE: Geometric control of cell life and death. *Science* 276:1425–1428, 1997
- Dascalu A, Mattheyou A, Oron Y, Korenstein R: A hyperosmotic stimulus elevates intracellular calcium and inhibits proliferation of a human keratinocyte cell line. *J Invest Dermatol* 117:714–718, 2001
- Görmär FE, Bernd A, Bereiter-Hahn J, Holzmann H: A new model of epidermal differentiation: Induction by mechanical stimulation. *Arch Dermatol Res* 282:22–32, 1990
- Kippenberger S, Bernd A, Loitsch S, Müller J, Guschel M, Kaufmann R: Cyclic stretch up-regulates proliferation and heat shock protein 90 expression in human melanocytes. *Pigment Cell Res* 12:246–251, 1999
- Kippenberger S, Loitsch S, Müller J, Guschel M, Ramirez-Bosca A, Kaufmann R, Bernd A: Melanocytes respond to mechanical stretch by activation of mitogen-activated protein kinases (MAPK). *Pigment Cell Res* 4:278–280, 2000a
- Kippenberger S, Bernd A, Loitsch S, Guschel M, Müller J, Bereiter-Hahn J, Kaufmann R: Signaling of mechanical stretch in human keratinocytes via MAP kinases. *J Invest Dermatol* 114:408–412, 2000b
- Takei T, Rivas-Gotz C, Delling CA, *et al*: Effect of strain on human keratinocytes in vitro. *J Cell Physiol* 173:64–72, 1997

Reply

To the Editor:

In reply to the remarks of Kippenberger *et al*, we would kindly refer the letter's authors to the ample work performed by Parsegian and coworkers, which demonstrates in detail that osmotic stress causes structural changes of macromolecules via crowding and hydration effects (Rand, 2000). For example, previous attempts to change the volume of ion channels by hydrostatic pressure were replaced by the elegant method of osmotic stress based on maintaining a difference in solute concentrations between the bathing solution and the water-filled ion channels, leading to decreased channel volume. Therefore, both mechanical pressure and osmotic stress cell membrane affect intracellular structures.

As to the other points, the periodical mechanical stimulation, employed by Görmär *et al* (1990), is the equivalent to a cyclic strain used by Takei *et al* (1997) and resulted in proliferative changes as reported. On the other hand, the technique of constant stretch gives different results as described by Kippenberger *et al*.

The presentation of skin stretch as a cause of "skin enlargement without changes in skin thickness" is an oversimplification of the clinical and pathologic skin conditions that were cited by Takei *et al* (1997). For example, the dermis might become thinner, and a consideration of the length of time of the mechanical stimulus application should be taken into account. Furthermore, it is difficult to draw conclusions using keratinocytes as the sole cellular model, without further evaluation of a fibroblast-keratinocyte 3D model, with a specific hormonal milieu relevant to the clinical comparison.

Cells respond to different mechanical stimuli by membrane deformation, to be followed by activation of specific signal transducers. Specifically, it should be noted that a hyperosmotic stress leads cell shrinkage, causing a fast increase in the ratio of cell surface to volume. The cellular deflation over a rigid skeleton leads to both membrane stretching and folding of the plasma membrane in various areas, thus mimicking mechanical stress. Additionally, the intracellular pressure mentioned by Kippenberger *et al* is not known as a major sensor in keratinocytes, and we believe the proliferative and differentiation effects are related to ion-channel activation and to interplay of calcium homeostasis and gradient, which are critical to the epidermis.

In conclusion, we delineated mechanical pathways of activation and their end-point results of proliferation-differentiation balance. The general study of volume control in keratinocytes, whose physiologic relevance is not clear to us, was not within the scope of our study.

Avi Dascalu, Yoram Oron, Rafi Korenstein
José Carlos García-Borrón
Sackler School of Medicine,
Tel-Aviv University, Ramat Aviv, Israel

REFERENCES

- Görmär FE, Bernd A, Bereiter-Hahn J, Holzmann H: A new model of epidermal differentiation: Induction by mechanical stimulation. *Arch Dermatol Res* 282:22–32, 1990
- Rand RP, Parsegian VA, Rau DC: Intracellular osmotic action. *Cellular Mol Life Sci* 57:1018–1032, 2000
- Takei T *et al*: Effect of strain on human keratinocytes in vitro. *J Cell Physiol* 173:64–72, 1997

Manuscript received March 12, 2001; accepted for publication March 12, 2001.

Reprint requests to: Dr. A. Dascalu, Department of Physiology and Pharmacology, Room 501, Sackler School of Medicine, Tel-Aviv University, Ramat Aviv 69978, Israel.